



DR. RAM MANOHAR LOHIA INSTITUTE OF MEDICAL SCIENCES
VIBHUTI KHAND, GOMTI NAGAR, LUCKNOW-10(U.P)
Phone No : - 0522-4918502, 4918510

Ref No: HRF/PAC/2025
1343

Dated: 17.Oct. 2025

Notice

Sub:- Procurement of following items on proprietary/ single quotation basis for different department

The Dr. RMLIMS, Lucknow intend to procure following item (s) manufactured as per mentioned against item name for **Mycobacterium w (Heat killed) Injection** on proprietary/ single quotation basis from their authorized dealer/ seller as per enclosed Technical Specifications.


Sl no	Product details	Principle Company (Manufacturer)	Subsidiary Company (Marketing)	Authorised Seller/ Company/Dealer
1	Mycobacterium w (Heat killed) Injection	M/s Cadila Pharmaceuticals Limited, 1389-Trasad Road, Ahmedabad (India) -387810	—	M/s Manglam Associates, Lucknow-226006.

The Proprietary Certificate for above items (s) submitted by principle company or their authorized seller/ Company/ Dealer is attached. The above documents are being uploaded for open information to all manufacturer/ supplier to submit comments / objections/ representation on the chairmen. (HRF), Dr RMLIMS, Lucknow on email id hrtendercell@gmail.com, from the date mentioned above, falling which it will be presumed that no other supplier is having any comment to offer and the case will be decided on merits. The comments/objection/representation to be submitted on the following:-

- Weather the above medicine/surgical items is manufactured by any other manufacture other than as per mentioned principle company or their Authorized seller/company/Dealer.
- Fulfill all the parameter(s) as per technical specifications.

Encl: - Related documents enclosed.

- (1) HRF Requisition form
- (2) PAC Certification of company letter.
- (3) Authorization from company letter.


Chairman (HRF)
Dr. RMLIMS, Lucknow

HRF Requisition Form

Request for new items/upgraded version (Drugs, Consumables & Disinfectants)

('X' if is not applicable)

1. Name of item (generic name only, no brand name)..... HEAT KILLED MICRO BACTERIUM (W) (MYCOBACTERIUM W)

Please note that if another brand of the same item is already available in HRF, the request will not be entertained for another brand.

2. Quantity needed (Per month) Quantity required
3. Probable Source (I) = 90 Injections
(II) for 30 patients
(III)
(If only one source please sign. The P-3 Form on back page)

4. Similar item available In HRF inventory? -Yes/No ✓
5. If yes then, why this item?

6. Do you want this item to be made available on regular basis -Yes/No ✓

7. If yes, then what will be the monthly consumption of this item?

8. Is same item in single unit be used on many patients ? If yes then specify the
Number of times/Number of patients, the unit will be used.....

9. Will it be a part of any procedure (Dossier, please specify the name of
procedure.....

10. If it is an upgraded version of an existing item in HRF inventory, do you want old
version to continue? - -Yes/No

11. Justification for new requisition..... For Gram negative

Septicaemia patients

Dr. Ishwar Ram Dhayal
MBBS, MS, M.Ch. (AIIMS)
Professor & Head
Dept. of Urology & Renal Transplant
Dr. R.M.L.I.M.S., Ganga Nagar, Lucknow
(Sign. of Consultant)

(Signature of Head of Department)

Dr. Ishwar Ram Dhayal
MBBS, MS, M.Ch. (AIIMS)
Professor & Head
Dept. of Urology & Renal Transplant
Dr. R.M.L.I.M.S., Ganga Nagar, Lucknow

Please note that new item will be processed for short-term rate contract this may take
about 1-2 months time.

Dr. Ram Manohar Lohia Institute of Medical Sciences, Lucknow

PROPRIETARY ARTICLE CERTIFICATE

It is certified that the items required should be purchased from M/s. MANGALAM ASSOCIATES Who are the sole manufacture/agents of the sole manufacturers M/s. CADILA PHARMA

Similar items manufactured by other firm(s) shall not be suitable for our purpose for the following reasons: -


- For Gram negative Septicaemia (where SOFA score > 6)
- No other company has this formulation.
- Injection required = 90 for 30 Patients

Requisition No:

Department :

Dated :

Urology
29/9/25


Dr. Ishwar Ram Dhayal
MBBS, MS, M.Ch. (AIIMS)
Professor & Head
Dept. of Urology & Renal Transplant
Dr. R.M.L.I.M.S., Gomti Nagar, Lucknow



Designation &
Sign of Head of
Department/ Section
Dr. Ishwar Ram Dhayal
MBBS, MS, M.Ch. (AIIMS)
Professor & Head
Dept. of Urology & Renal Transplant
Dr. R.M.L.I.M.S., Gomti Nagar, Lucknow

N.B. The Indenter before recording the above certificate should satisfy himself that the article is genuinely of proprietary nature manufactured under patent laws.



Food & Drugs Control Administration

BLOCK NO. 8, 1ST FLOOR, DR. JIVRAJ MEHTA BHAVAN,
GANDHINAGAR, GUJARAT STATE, INDIA. PIN : 382010



Certificate No. : **22113688**

On the basis of the inspection carried out on **09-10/11/2022** we certify that the site indicated on this certificate complies with Good Manufacturing Practices for the dosage forms, categories and activities listed in Table 1.

1 Name & Address of site : **CADILA PHARMACEUTICALS LIMITED**
1389, TRASAD ROAD, DHOLKA - 382 225
DISTRICT : AHMEDABAD,
GUJARAT, INDIA

2 Manufacturer's Licence number : **G/1500** **G/1090**

3 Table : 1

Dosage Form (s)	Category (ies)	Activity (ies)
Tablet, Capsule, Oral Liquid, Small Volume Parenteral (Liquid, Ophthalmic Preparation - Vials & Ampoules) & Lyophilized Injection (Vials), Oral Powder (Granules) and Bulk Drug (API- Streptokinase)	General	Manufacturer
Tablet, Capsule, Dry Syrup, Small Volume Parenteral (Dry Powder - Injection)	Beta-lactum	
Tablet, Capsule, Dry Syrup, Small Volume Parenteral (Dry Powder - Injection)	Cephalosporin	

The responsibility for the quality of the individual batches of the pharmaceutical products manufactured through this process lies with the manufacturer.

This certificate remains valid until **16/11/2025** It becomes invalid if the activities and /or categories certified herewith are changed or if the site is no longer considered to be in compliance with GMP

Format of this certificate is as per WHO TRS No. 908 of 2008.

Address of certifying authority

Food & Drugs Control Administration, Block No. 8, 1ST floor, Dr. Jivraj Mehta Bhavan, Gandhinagar, Gujarat State, India. - Pin : 382010



Name & function of : **(Dr. H. G. KOSHIA)**
responsible Person **Commissioner**

Email : **comfdca@gujarat.gov.in**

Phone : **91-79-23253417**, Fax : **91-79-232-53400**

Date : **17/11/2022**

CERTIFICATE OF ANALYSIS
QUALITY CONTROL DEPARTMENT

Product Name	:	SEPSIVAC		
Generic Name	:	MYCOBACTERIUM W (HEAT KILLED) INJECTION		
Batch Size	:	5000 Vials	A.R. Number	: FG/DHLK/2404007
Batch Number	:	GO24002	Pack Style	: Vial
Mfg. Date	:	Sep. 2024	Exp. Date	: Aug. 2027
Specification Number	:	DP-A797-014-04	STP No.	: STP-A797-014-04
Date of Analysis	:	16/09/2024	Date of Report	: 12/10/2024
S.No.	TESTS	RESULTS	SPECIFICATION	
1	Description	Colorless opaque suspension in which cells are suspended having tendency to settle on keeping aside	Colorless opaque suspension in which cells are suspended having tendency to settle on keeping aside	
2	Identification	Acid fast Mycobacterium identified by Zeil-Neilson staining	Should comply for identification of acid fast Mycobacterium by Zeil-Neilson staining	
3	pH	7.1	6.0 to 7.5	
4	Sterility	Sterile	Should be sterile	
5	Bacterial Endotoxin test	Less than 25 EU/ml	Not more than 25 EU/ml	
6	Extractable Volume	0.70 ml	Not less than 0.6 ml	
7	Light absorbance	0.80	0.6 to 0.9	
8	Sodium chloride content (0.9% w/v)	0.97 % w/v	0.81%w/v to 0.99%w/v	
9	Thiomersal	0.010 % w/v	0.008% w/v to 0.012% w/v	
10	Assay			
	Animal Assay (Nodule test)	Greater than and equal to 2.3 mm	Not less than 1.5 mm	

Conclusion: The Product is standard quality in respect of above specification.

Prepared By :

Date :

(Executive-Q.C.):

Checked By :

Date :

(Manager-Q.C.):

Approved By :

Date :

(Head-Q.C.):

FORMAT NO. : FQC214-02/00

FORM 26

(See Rules 73 and 83)

Certificate of Renewal of Licence to manufacture for sale of Drugs Other than those specified in Schedule X

Certified that Licence in ~~Form 25~~ No:G/1500 Granted on the Date: 06/07/1999
Form 28 No:G/1090 Granted on the Date: 06/07/1999

To M/s. CADILA PHARMACEUTICALS LTD.

for the manufacture of the following categories of drugs being drugs other than those specified in Schedule C and C (1) and "X" to the Drugs and Cosmetics Rules, 1945 drugs Specified in Schedule C and C (1) excluding those specified in Schedule "X" to the Drugs and Cosmetics Rules, 1945,

at the premises situated at PLOT NO. 1389, TRASAD ROAD, DHOLKA, DIST :
AHMEDABAD

has been renewed from : 01/01/2018 To 31/12/2022

Name(s) of drugs (each item to be separately specified) : As per list Approved & Annexed

Names of approved competent technical staff

: As per list Approved & Annexed.



Signature :

Designation :

Commissioner
Food & Drugs Control Administration
Gujarat State.

Date:- 07/02/2018



BY REGD.A.D.

NO.MFG/CADILA/16783/B
COMMISSIONER of,
Food & Drugs Control Administration,
Block No.8, 1st floor, Dr. Jivraj Mehta Bhavan,
GANDHINAGAR. - 382 010.

DATE :

14 FEB 2018

To,

CADILA PHARMACEUTICALS LTD.,
PLOT NO. 1389, TRASAD ROAD, DHOLKA,
DIST : AHMEDABAD

SUB : Renewal of licence for the Period : 2018 - 2022

Sir,

REF : Your application in Form No. 24 & 27 Dated : 30/12/2017

Licence having following details has been renewed for a period of Five Years.

I. Details of Licence :

II.

Form No.	Licence No.	Date of Licence	Renewed for a Period		Category
			From	To	
25	G/1500	06/07/1999	01/01/2018	31/12/2022	TABLET, GRANULES, CAPSULE, ORAL LIQUID, POWDER
28	G/1090	06/07/1999	01/01/2018	31/12/2022	TABLET, GRANULES, CAPSULE, ORAL LIQUID, SVP, DRY SYRUP, BULK DRUG

II. Approved Technical Person (Manufacturing Section)

1. AS PER LIST APPROVED AND ANNEXED

Approved Technical Person (Testing Section)

1. AS PER LIST APPROVED AND ANNEXED

Further you are also permitted to get your raw materials and finished products tested at any govt. approved public testing laboratories holding approval in form 37 under Drugs & Cosmetics Rules 1945, only for those tests which require sophisticated equipments.

You are requested to get the products amended as per latest I.P. wherever applicable.

Further you are asked for compliance with respect to DCGI notification dated 07/08/2014 for products containing single ingredients (except for export purpose)

Yours faithfully,

For Commissioner,

Food & Drugs Control Administration,

Encl:- 1. Renewal Certificate

2. List of Products Approved

NO.MFG/CADILA/

/B.D. :-

Copy with a copy of approved list of products forwarded to :-

1) The Assistant Commissioner, AHMEDABAD (RURAL), GANDHINAGAR

2) The Statistical Cell, (Ct. Branch)

For Commissioner,

Food & Drugs Control Administration.

RETENTION OF LICENCE

Retention of Licence to manufacture for sale (or for distribution) of drugs specified in Schedules C, C(1) excluding those specified in schedule X

Certified that Licence in Form 28 No: G/28/1090 Issue Date: 06-Jul-1999

To M/s. CADILA PHARMACEUTICALS LIMITED

to manufacture the following categories of drugs being drugs Specified in Schedules C and C (1) excluding those specified in Schedule "X " to the Drugs and Cosmetics Rules, 1945, at the premises situated at 1389, TRASAD ROAD, DHOLKA-382 225, DISTRICT:AHMEDABAD, GUJARAT, INDIA

has been retained from: 01-Jan-2023 To 31-Dec-2027

Name(s) of drugs : As per list Approved

Names of approved competent technical staff : As per list Approved

Signature :

(This Document is Digitally Signed.)
Dr. H. G. KOSHIA

Designation :

Commissioner
Food & Drugs Control Administration
Gujarat State

Date: 16-Dec-2022

Note: You are requested to apply for the Retention of the above licence in 3 months before its VALIDITY EXPIRES.



Reg ID : 631706

Doc ID: RC945700001090

CADILA PHARMACEUTICALS LIMITED

Print Date : 16/12/2022 11:23 AM

Signature valid

Digitally signed by
Date: 2022.12.16
11:20:42 +05:30
Reason: Sign
Location: AHD

RETENTION OF LICENCE

Retention of Licence to manufacture for sale (or for distribution) of drugs other than those specified in Schedules C, C (1) and X

Certified that Licence in Form 25 No: G/25/1500 Issue Date: 31-Dec-2012

To M/s. CADILA PHARMACEUTICALS LIMITED

to manufacture the following categories of drugs being drugs other than those specified in Schedules C and C (1) and Schedule "X" to the Drugs and Cosmetics Rules 1945, on the premises situated at 1389, TRASAD ROAD, DHOLKA- 382225, DISTRICT:AHMEDABAD, GUJARAT, INDIA

has been retained from: 01-Jan-2023 To 31-Dec-2027

Name(s) of drugs

: As per list Approved

Names of approved competent technical staff

: As per list Approved

Signature :


(This Document is Digitally Signed.)
Dr. H. G. KOSHIA

Designation :

Commissioner
Food & Drugs Control Administration
Gujarat State

Date: 16-Dec-2022

Note: You are requested to apply for the Retention of the above licence in 3 months before its VALIDITY EXPIRES.



Reg ID : 631699

Doc ID: RC077300001500

CADILA PHARMACEUTICALS LIMITED

Print Date : 16/12/2022 11:21 AM

Signature valid

Digitally signed by
Date: 2022.12.16
11:24:27 +05:30
Reason: Sign
Location: AHD

in both 0.2 and 0.3 groups ($p=0.0001$). SOFA scoring clearly showed that patients receiving 0.3 *Mycobacterium w* (Heat Killed) injection had earliest decrease in scores with 83% of the patients recovering on day 15 and 100% on day 22 as compared to other groups receiving lesser doses as well as the standard treatment group. Patients receiving 0.3 *Mycobacterium w* (Heat Killed) injection dose had significant early recovery of organ function including renal and respiratory, multisystem organ failure, fever, and there was an early improvement of SOFA score. Patients receiving 0.2 *Mycobacterium w* (Heat Killed) injection dose also had statistically significant improvement in early recovery in hepatic, respiratory, cardiovascular, multisystem organ failure and fever. Patients receiving 0.1 mL dose had marginal early recovery as compared to the standard treatment group although it was not statistically significant except in case of fever. Mild local reactions were seen in 2 patients in 0.1 group, 3 patients in 0.2 group and 2 patients in 0.3 group. There was no serious adverse events observed. Cytokine levels were significantly elevated at baseline in all the groups which was more significant fall in 0.3 group compared to other treatment group. It was concluded that 0.3 *Mycobacterium w* (Heat Killed) injection dose seems to be efficacious in early organ function recovery and microbiological resolution among all the groups studied. Moreover, *Mycobacterium w* (Heat Killed) injection may serve as a promising novel immunomodulatory therapy in improving outcome in severe Gram Negative Septic Shock.

Study 2 - Patients were randomized 1:1 to the experimental (*Mycobacterium w* (Heat Killed) with standard therapy) or the control arm (standard therapy). Fifty patients with severe sepsis (25 *Mycobacterium w* (Heat Killed), 25 control) were included in the study. There were 7 and 8 deaths in the *Mycobacterium w* and control groups, respectively ($p=0.51$). The days on mechanical ventilator were significantly lesser in the *Mycobacterium w* (Heat Killed) group compared with control (median, 6 vs 9; $p=0.025$). The median ICU and hospital length of stay was significantly less in the *Mycobacterium w* (Heat Killed) arm (7 vs 12 days [$p=0.006$] and 10 vs 16 [$p=0.007$], respectively). The delta SOFA score was significantly higher in the control arm ($p=0.027$). There was a higher incidence of secondary bacterial infections in the control group ($P=0.009$). Therefore, the use of *Mycobacterium w* (Heat Killed) in severe sepsis was associated with significant reduction in days on mechanical ventilation, ICU and hospital length of stay, lower incidence of nosocomial infection, and delta SOFA score.

Study 3 - Patients were randomized 1:1 to the experimental (*Mycobacterium w* (Heat Killed) with standard therapy) or the control arm (standard therapy). One hundred one patients with severe sepsis due to Gram Negative Bacteria (101 *Mycobacterium w* (Heat Killed), 101 control) were included in the study. There were 8 and 18 deaths in the *Mycobacterium w* (Heat Killed) and control groups, respectively. The primary end point of the study (28 days mortality) using *Mycobacterium w* (Heat Killed) as an adjuvant to standard treatment in sepsis to reduce the 28 days mortality associated with sepsis was met. The difference between the two treatment arms was 10.89% (absolute reduction) and 55.56% (relative reduction) which is clinically relevant and statistically significant. ($p=0.029$). There was significant difference in time to discharge ($p=0.0001$) and time to discontinuation of vasopressor treatment ($p=0.0159$) between the two arms, while no significant difference was found in ICU days and Ventilator days between the two arms. The observed difference is probably due to higher and early mortality in the control arm. The reduction in mean SOFA score from baseline to day 28 is observed in both arm. However, the reduction in SOFA score is significantly more in Treatment Arm as compared to Control Arm ($p=0.0182$ at day 14, $p=0.0079$ at day 21 and $p=0.0181$ at day 28). Therefore, the use of *Mycobacterium w* (Heat Killed) in severe sepsis was associated with significant reduction in days on mechanical ventilation, ICU and hospital length of stay, lower incidence of SOFA score.

5.3 Pharmacokinetic properties

Not applicable

6. Non Clinical Properties

6.1 Animal Toxicology or Pharmacology:

Non-clinical data reveals no special hazard for human based on conventional studies like acute toxicity studies in mice & rats and subacute toxicity studies in mice & rabbits.

7. Pharmaceutical particulars

7.1 Incompatibilities.

Mycobacterium w (Heat Killed) Injection is not recommended to be diluted with any i.v. fluids.

7.2 Shelf life

The shelf life of *Mw* (Heat Killed) injection is 36 months when stored at +2°C to +8°C.

7.3 Packaging information

0.6 mL of *Mycobacterium w* (Heat Killed) Injection is presented in USP type 1 glass vial.

7.4 Storage and handling instructions

Store at +2°C to +8°C.
Do not freeze.
Discard if frozen.
Shake well before use.
Keep out of reach of children.
Protect from light.
Do not use *Mw* (Heat Killed) injection after the expiration date shown on the label.

8. Patient Counselling Information.

Consult the physician for personalized medical advice

9. Manufacturer:



Cadila Pharmaceuticals Limited
1389, Trasad Road,
Dholka, District - Ahmedabad, Gujarat
Phone: +91-952714-22148/183/84
Fax: +91-2714-220315, +91-2714-221848
Website: www.cadilapharma.com

10. Market authorization with date

MF/BI/O/19/09/057 dated 28-Nov-2019

11. Last Revision details:

December 2019

For the use only of a registered medical practitioner or a Hospital or a Laboratory

***Mycobacterium w* (Heat Killed) Injection**

Sepsivac

1. Description:

Mycobacterium w (Heat Killed) Injection is a colourless opaque suspension in which *Mycobacterium w* cells are suspended and having tendency to settle down during storage.

2. Qualitative and Quantitative composition:

Each 0.1 mL contains:	
<i>Mycobacterium w</i> (Heat Killed)	: 0.5 x 10 ⁸ Cells
Sodium Chloride IP	: 0.9 % w/v
Thiomersal IP	: 0.01 % w/v
Water for injections IP	: q.s

3. Dosage form and strength:

Mycobacterium w (Heat Killed) injection is a sterile suspension for intradermal injection. The strength of *Mycobacterium w* (Heat Killed) Injection is 0.5 x 10⁸ Cells per 0.1 mL.

4. Clinical particulars:

3.1 Therapeutic Indication:

Mycobacterium w (Heat Killed) injection is used as immunotherapeutic agent in the following disease conditions:

1. Leprosy – in Lepromin negative patients
2. Advanced Non-Small Cell Lung Cancer (NSCLC) - in combination with Paclitaxel plus Cisplatin regimen
3. Sepsis (due to gram Negative infections) - as adjuvant to the standard treatment

3.2 Posology and method of administration :

Posology:

1. Leprosy

The first dose of *Mycobacterium w* (Heat Killed) injection is administered by two intradermal injections, i.e. each 0.1 mL administered at left and right hand of deltoid regions. Further, 0.1 mL of *Mw* (Heat Killed) injection is administered in one deltoid area every 3 months interval for 2 years or advised by physician. After initial dose, total of 8 doses is recommended to be taken at 3 monthly interval along with regular multi drug therapy (MDT).

2. Non-Small cell lung cancer (NSCLC)

Adult and Geriatric patients (18 years and above)

It is administered intradermally. It is advisable to inject 0.1 mL of the drug per site. The amount of the drug to be administered at one time and its frequency of administration are described to be dependent on therapeutic condition.

Usually 0.2 mL of *Mycobacterium w* (Heat Killed) injection is recommended to be given initially in two-divided dose of 0.1 mL on each arm followed by 0.1 mL subsequently. The initial dose of 0.2 mL will be given prior to one week of onset of chemotherapy and subsequent dose of 0.1 mL each will be administered in adjuvant to four cycles of chemotherapy (Cisplatin and Paclitaxel) including 3 week (21 days) each cycle. *Mw* (Heat Killed) injection is recommended to be taken at 2nd and 3rd week of each cycle (Intradermal 0.1 mL in deltoid).

Mw (Heat Killed) injection should be administered monthly after completion of all chemotherapy cycles (four cycle) till 12 months from start of treatment.

3. Severe Sepsis:

A 0.3 mL of *Mycobacterium w* (Heat Killed) injection is given intradermally in three divided doses of 0.1 mL each on three different sites daily for three days. Total dose is 0.9 mL of *Mycobacterium w* over a period of three days. *Mw* (Heat Killed) injection is recommended to be prescribed in patient with age 18-65 years.

Method of administration

The recommended site for giving the *Mw* (Heat Killed) injection is at the insertion of the deltoid muscle near the middle of the left upper arm. Sites higher on the arm are more likely to lead to keloid formation, the tip of the shoulder particularly. For cosmetic reasons, a scar on the upper and lateral surface of the thigh may be preferred and this is an alternative site.

The upper arm must be approximately 45 degrees to the body. This can be achieved if the hand is placed on the hip with the arm abducted from the body. The skin should be swabbed with spirit and allowed to dry. It is advisable to use 26 G or smaller gauge (27 G, 30 G) needle. The operator stretches the skin between the thumb and forefinger of one hand and with the other slowly inserts the needle, with the bevel upwards, till bevel is fully in the dermis and not visible out. The needle can usually be seen through the epidermis. A correctly given intradermal injection results in a tense blanched raised bleb (Peau D'Orange) and considerable resistance is felt when the fluid is being injected.

A bleb typically of 7 mm diameter follows a 0.1 mL injection. If little resistance is felt when injecting and a diffuse swelling occurs as opposed to a tense blanched bleb, the needle is too deep. The needle should be withdrawn and reinserted intradermally before more is injected. The subject must always be advised of the normal reaction to the injection. The second dose of injection is to be given one inch apart from previous dose to minimize the chance of local reaction.

This injection must be given strictly intradermally.

Injection site reaction and care of the injection site:

Following intradermal administration of *Mw* (Heat Killed) injection, normally a local reaction develops at the immunization site within two to six weeks, beginning as a small papule which increase in size for a few weeks widening into a circular area with scaling, crusting and occasional bruising. Occasionally a shallow ulcer develops. It is not necessary to protect the site from becoming wet during washing and bathing, but should any oozing occur, a temporary dry dressing may be used until a scab forms. It is essential that air be not excluded. If absolutely essential an impervious dressing may be applied but only for a short period (for example, to permit swimming) as it may delay healing and cause a larger scar. The lesion slowly subsides over several months and eventually heals leaving only a small, flat scar.

3.3 Contraindications:

- Mycobacterium w (Heat Killed) Injection** is contraindicated in
 - History of allergic reactions attributed to **Mycobacterium w (Heat Killed) Injection** or any of the excipients in the formulation mentioned in section 2
 - Individuals with fever
 - Pregnant and lactating women
 - Individuals with generalized septic skin conditions (if eczema exists, a site should be chosen that is free from skin lesions)
 - Patient with chronic debilitating condition other than the proposed indication

3.4 Special warnings and precautions for use:

- Injection site reaction and care of the injection site
- Please see section 3.2 (posology and method of administration) for details information

Severe injection site reactions, large ulcers and abscesses are most commonly caused by faulty injection technique. Hence adequate precaution should be taken while intradermal injections and should be administered by a healthcare staff well trained in the procedure.

Mycobacterium w (Heat Killed) Injection can cause erythema, induration and ulceration of the skin at site of injection which are usually mild and can be self-healing. If the condition is not cured then please consider suspending therapy and necessary antibiotics.

Mycobacterium w (Heat Killed) Injection is not recommended to be administer via Intravenous, subcutaneous, intramuscular injection.

The intradermal injection can also cause minor active local reactions and local delayed type hypersensitivity reaction, type I and II reaction, neuritis. If condition get worsened or not self-cured please provide patient with proper supporting drug therapy.

3.5 Drug Interactions:

Mycobacterium w (Heat Killed) Injection is not known to have any drug-drug interactions. It is able to induce body's own immune response even when administered with other drugs like cytotoxic, anti-leprosy drugs.

3.6 Use in special populations:

Animal reproduction studies have not been conducted with **Mycobacterium w (Heat Killed) Injection**. It is also not known whether **Mycobacterium w (Heat Killed) Injection** can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity.

It is not known whether drug is excreted in human milk. As many drugs are excreted in human milk, caution should be exercised when drug is administered to a nursing woman.

3.7 Effects on ability to drive and use machines:

No studies on the effects on the ability to drive and use machines have been performed with **Mycobacterium w (Heat Killed) Injection**.

3.8 Undesirable effects:

It has been found to be generally well tolerated and free from severe systemic adverse effects in Leprosy treatment. The only side effects encountered were injection site erythema and ulceration. The erythema appeared after 48 hours of injection and was followed by induration by 7th day culminating into the formation of a shallow, self-healing ulcer in the 3rd week which healed with scar formation in the 4th week leaving a scar, as observed in clinical trials.

Severe injection site reactions, large ulcers and abscesses are most commonly caused by faulty injection technique where part or the entire dose is administered too deeply.

Keloid formation at the injection site is an uncommon and largely avoidable, complication of **Mw (Heat Killed) Injection**. Some sites are more prone to keloid formation than others and those using **Mw (Heat Killed) Injection** should adhere to the two sites recommended. (The mid-upper arm or the thigh). Most experience has been gained in the use of the upper arm and it is known that the risk of keloid formation is increased manifold when the injection is given at a site higher than the insertion of the deltoid muscle near the middle of the upper arm.

Clinical trials conducted in patients with NSCLC has shown anaemia, Leukopenia, Neutropenia, as treatment-emergent haematological toxicities (**Mycobacterium w (Heat Killed) Injection** with paclitaxel and cisplatin or paclitaxel and cisplatin). Out of which, anaemia was the most commonly reported. There were no grade 3 neutropenia cases reported in test arm whereas 2 cases of neutropenia (**Mycobacterium w (Heat Killed) Injection** with paclitaxel and cisplatin) were reported in control arm (paclitaxel and cisplatin). The overall incidences of grade 3 and 4 non-haematological laboratory toxicities were low in both arms. Among the two arms, the most common grade 3 and 4 non-haematological toxicities were increased transaminase and alkaline phosphatase levels. Clinical Adverse Events includes nausea and vomiting, diarrhoea, pain, weakness, weight loss, anorexia, neuropathy, Constipation, Breathlessness, Loss of appetite, Cough, Haemoptysis, Reactogenic reactions, Neuropathy and alopecia were observed in test arm. Out of these, Reactogenic reactions (96 out of total 277 adverse events), Nausea and Vomiting (56 out of total 277 adverse events) were most common. All these adverse events were grade 1 and 2. Only 2 patients experienced grade 3 (Diarrhoea and cough) adverse events.

In randomized, double-blind, comparative study conducted in patient with sepsis, the Mw group had significantly lesser incidence of secondary bacterial infection compared with the control group. Ventilator associated pneumonia (VAP) was the commonest secondary bacterial infection in both groups; 12 patients in the control group developed VAP, whereas 5 patients in the Mw group had occurrence of VAP. *Acinetobacter baumannii* was the commonest organism responsible for nosocomial infections. Catheter-related blood stream infection (CRBSI) occurred in 2 and 6 cases in the Mw and control groups, respectively. Six patients had both VAP and CRBSI: 2 in the Mw group and 4 in the control arm. In another Clinical Trial (Phase IIa) conducted in patient with sepsis a total of 30 out of 72 randomized patients reported adverse events which includes transfusion associated reaction, Thrombocytopenia, Hypokalemia, Transaminitis, Local site reaction, Hyponatremia, Hypokalemia, Respiratory system involvement, Deranged RFT. All the reported adverse events were mild and none were severe. Local injection site reactions were found to be probably related with the study drug. Transaminitis and Thrombocytopenia were possibly related with study drug (standard treatment and 0.1 ml of MW injection). In another clinical trial (phase IIb) conducted in patient with severe sepsis, twenty seven (27) SAEs were observed. Out of these, eight (8) SAEs were observed in Test treatment arm and nineteen (19) SAEs were observed in Control arm. This indicates **Mycobacterium w (Heat Killed) Injection** is found to be safe and well tolerated, without any major safety concerns in patients with severe sepsis.

3.9 Overdose:

No data is available on overdose with **Mycobacterium w (Heat Killed) Injection**.

5. Pharmacological properties

5.1 Mechanism of Action:

Immunomodulators or biological response modifiers have been in vogue since long. They work by altering biological response of individuals to various pathological conditions/disease. **Mycobacterium w (Heat Killed) Injection** is one of

such agent. It contains heat killed **Mycobacterium w** as an active drug substance. It elicits potent cell mediated immune response when administered intradermally. Cell mediated immune response are designated as the Th1 type or Th2 type depending on the cytokines liberated by excited T cells. Interferon Gamma & Interleukin - 2 are and Th2 response. Certain diseases are associated with decreased Th1 response. They include Malignancy and Leprosy. Improving Th1 response in such conditions is associated with improved outcomes.

5.2 Pharmacodynamics Properties:

Mycobacterium w (Heat Killed) shares antigens with *M. Tuberculosis* as well as **Mycobacterium Lepae**. In experimental models it is found to induce lympho-proliferative response. The lymphoproliferative response induced by **Mycobacterium w (Heat Killed)** is the most potent of all known immune modulators. It provides immunity against tuberculosis in animals. The protection provided against tuberculosis by **Mycobacterium w (Heat Killed)** is universal. It is seen in all species & strains. BCG also provides protection in animals. However, protection provided by BCG is species and strain specific and not universal. **Mycobacterium w (Heat Killed) Injection** provides protection against tuberculosis in species and strains of animals in which BCG works as well as in species and strains in which BCG fails to provide protection.

Cell mediated immune response are designated as the Th1 type or Th2 type depending on the cytokines liberated by excited T cells. Interferon Gamma & Interleukin - 2 are cytokines associated with Th1 response. In normal healthy individuals there is a harmony between Th1 response and Th2 response. Administration of **Mycobacterium w (Heat Killed)** is also associated with release of Th1 type cytokines like Interferon Gamma and Interleukin - 2.

Clinical Studies in LEPROSY

Mycobacterium w (Heat Killed) Injection exhibits an immune stimulant effect as shown by conversion of lepromin negative reaction into positive. In a clinical study of lepromin-negative BB (Borderline) leprosy patients, 75% patients became lepromin positive after the first dose, 81 % after the second dose and 100% after the third dose. In BL (Borderline Lepromatous) leprosy the conversion rates were 43%, 60% and 82% after the first, second, and third doses of **Mycobacterium w (Heat Killed) Injection**, respectively.

Lepromin conversion in response to **Mycobacterium w (Heat Killed) Injection** is attributed to the presence of the "right mix" of antigenic determinants in the preparation which stimulate cell mediated immunity (CMI). This leads to histopathological upgrading and clearance of dermal granuloma without the risk of hypersensitivity reactions which could lead to nerve damage.

Clinical trials have revealed the beneficial effects of Immunomodulation with heat killed **Mycobacterium w (Heat Killed) Injection** in accelerating the bacillary clearance and shortening the duration of therapy in multibacillary leprosy. It has been shown to produce statistically significant ($p < 0.001$) bacteriological fall in LL and BL leprosy from 6 months onwards.

Histopathologically, the addition of **Mycobacterium w (Heat Killed) Injection** to MDT (multi-drug therapy), has shown a higher and statistically significant upgradation in LL and BL leprosy ($P < 0.001$) and complete clearance of granuloma as compared to patients treated only with MDT. Therefore, **Mycobacterium w (Heat Killed) Injection** is recommended as an adjunct to MDT in multibacillary leprosy to induce an early bacteriological clearance and shorten the duration of therapy.

Clinical Studies in NSCLC

Study 1 - Between January 2005 and June 2005, patients with advanced NSCLC were randomized to 2 arms. In arm A, 26 patients were randomized to receive Cisplatin 75 mg/m² on Day 1, and Etoposide 100 mg/m² on Days 1-3. The cycle was repeated every 21 days. **Mycobacterium w (Heat Killed) Injection** 0.1 mL was administered intradermally every 14th day for 6 months. In arm B, 22 patients were randomized to Cisplatin 75 mg/m² on Day 1 and Etoposide 100 mg/m² on Days 1-3. The cycle was repeated every 21 days.

Mycobacterium w (Heat Killed) Injection improved response rate of chemotherapy by 11% (38% in **Mycobacterium w (Heat Killed) Injection** arm vs 27% in control arm). **Mycobacterium w (Heat Killed) Injection** also improved median survival by 4 months (11 months in **Mycobacterium w (Heat Killed) Injection** arm vs 7 months in control arm).

The improvement in response rate and improved median survival seen in **Mycobacterium w (Heat Killed) Injection** arm were associated with lower incidence of hematological and non-hematological toxicities as compared to the control arm.

Use of **Mycobacterium w (Heat Killed) Injection** improves response rate and median survival in NSCLC.

Study 2 - Patients with advanced NSCLC were randomized to 2 groups. The treatment group (n=9) received chemotherapy (cisplatin-etoposide) and radiotherapy. **Mycobacterium w (Heat Killed) Injection** was administered intradermally. The control group (n=10) received chemotherapy (cisplatin-etoposide) and radiotherapy. The **Mycobacterium w (Heat Killed) Injection** group had better tolerance to therapy (interruption of therapy in 50 % vs 0%). At the end of therapy 67 % responded to therapy in **Mycobacterium w (Heat Killed) Injection** arm compared to none in control arm. The lung function improved in all the 9 patients in **Mycobacterium w (Heat Killed) Injection** arm. The response seen in **Mycobacterium w (Heat Killed) Injection** arm were durable in nature as seen to be stable during follow up at (6 months or more).

Study 3 - Two hundred and twenty-one treatment naïve patients with stage IIIB and IV NSCLC were randomized to receive paclitaxel and cisplatin with or without **Mycobacterium w (Heat Killed) Injection** administered intradermally, 0.1 mL on each deltoid on the first visit at least 1 week prior to 1st cycle and then on day 8 and 15 of each cycle of chemotherapy. The cycles were repeated every 3 weeks for a total of 4 cycles. **Mycobacterium w (Heat Killed) Injection** was administered once a month thereafter until progression or for a maximum of 12 months from the start of treatment in the investigational arm. The primary endpoint of the study was overall survival (OS) and the secondary outcome measures were response rate (RR) and progression free survival (PFS). One hundred twelve pts were randomized to the control arm and 109 pts to the **Mycobacterium w (Heat Killed) Injection** arm.

Mycobacterium w (Heat Killed) Injection improved response rate by 11% (36% in the control arm vs. 47% in the **Mycobacterium w (Heat Killed) Injection** arm). There were three complete responses, all in the **Mycobacterium w (Heat Killed) Injection** arm. Use of **Mycobacterium w (Heat Killed) Injection** was associated with improvement in survival also. Improvement in Median PFS with use of **Mycobacterium w (Heat Killed) Injection** was 96 days [$p = 0.0446$; HR 0.43 (95% CI 0.25-0.73)] from 157 days in control arm to 253 days in **Mycobacterium w (Heat Killed) Injection** arm. Use of **Mycobacterium w (Heat Killed) Injection** also improved median survival by 59 days [$p = 0.0034$; HR 0.55 (95% CI 0.37-0.82)] from 236 days in control arm to 295 days in the **Mycobacterium w (Heat Killed) Injection** arm.

The improvement in response rate and survival seen with **Mycobacterium w (Heat Killed) Injection** were associated with adverse events comparable to the control arm.

Use of **Mycobacterium w (Heat Killed) Injection** in patients with advanced NSCLC is safe and results in both improvement in progression free survival and overall median survival.

Clinical studies in SEVERE SEPSIS

Study 1 - Phase IIa study with 72 patients having gram negative severe sepsis/septic shock were enrolled in 3 groups (escalating dose levels 0.1, 0.2 and 0.3 **Mycobacterium w (Heat Killed) Injection** over initial 3 days with standard therapy versus standard therapy alone). Significant early clinical and microbiological resolution was seen

F. No. BIO/MA/19/000020
Government of India
Directorate General of Health Services
Central Drugs Standard Control Organization
(Biological Division)

FDA Bhawan, Kotla Road,
New Delhi- 110002.
Dated:

To

M/s Cadila Pharmaceuticals Limited,
1389-Trasad Road, Dholka,
Ahmadabad (India) – 387810.

02 MAR 2020

Subject: Correction in the Permission to Manufacture *Mycobacterium w* (Heat Killed) injection in Form 46 - regarding.

References:

1. Your application no. CPL/RA/21-01-1219 dated 21.12.2019 submitted to this office vide diary no 17098 dated 24.12.2019
2. Manufacturing permission in Form 46 issued by this office through SUGAM online portal on 04.12.2019

Sir,

In continuation to Manufacturing Permission in Form 46 issued vide permission no. MF/BIO/19/000057, dated 28.11.2019 and based on the documents submitted by you, this Directorate has no objection for amendment in said permission as mentioned below:

1. The name of the New Drug to be read as "*Mycobacterium w* (Heat Killed) Injection" instead of "*Mycobacterium w* (Heat Killed) Vaccine"
2. The presentation under "Dosage Form" to be read as "0.6 ml in 2 ml glass vial (USP Type I)" instead of "0.5 ml glass vial (USP Type I)".
3. Indication to be read as "Indicated as an adjuvant to standard treatment in sepsis (due to gram negative infections) when given as 0.3 ml in three divided doses of 0.1 ml each on three different sites, daily for three days in age group of 18 to 65 years" instead of "Indicated as an adjuvant in sepsis (due to gram negative infections) treatment when given as 0.1ml for three consecutive days in age group of 18 to 65 years"

Further, you are requested to submit PMS study protocol for approval as mentioned in the condition no 2 in the cover letter of said permission.

However, all other terms & conditions as stipulated in the said permission issued by this Directorate shall remain unchanged.

Yours faithfully,



(Dr. V. G. Somani)
Drugs Controller General (India)

An Adjunct to Standard Treatment in
Sepsis due to gram negative infections


Sepsivac[®]
(Heat killed Mw)

Save More Lives




India's Novel Immunomodulator

Mean SOFA Score Over Time



Sepsivac led to significant reduction of SOFA score.



Each kit contains:

- 1 vial ■ 3 syringes
- 3 needles of 24 G (to draw medicine)
- 3 needles of 26 G (to inject medicine)

ABM Name :

Mob. No.:

Reference: 1. Reference: Sepsis Alliance (2022). Available at: <https://www.sepsis.org/sepsis-basics/what-is-sepsis> | 2. J Thorac Dis. 2017 Feb;9(2):392-405. doi: 10.21037/jtd.2017.02.10 | 3. Data On File | 3. Sehgal IS, Basumatary NM, Dhooria S, Prasad KT, Muthu V, et al. A Randomized Trial of Mycobacterium w in Severe Presumed Gram-Negative Sepsis. Chest. 2021;160(4):1282-1291

Care
Cadila VoltaCare
A Division of Cadila Pharmaceuticals Ltd.

CADILA
PHARMACEUTICALS
LIMITED
The Care Continues...

For more info...